

ANGLE MEASURES OF SOME CONES ASSOCIATED WITH FINITE REFLECTION GROUPS

P. V. Bibikov¹, V. S. Zhgoon²

Let V be a Euclidean vector space of dimension n with the inner product $(\cdot, \cdot)$. For a convex polyhedral cone C we denote by C° its open kernel, by $\langle C \rangle$ its linear span, and by C^* its *dual cone*, i.e. $C^* = \{v \in V : (v, u) \geq 0 \ \forall u \in C\}$. Denote by $\sigma(C)$ the *relative angle measure* of C , i.e. $\sigma(C) = \text{vol}(C \cap B) / \text{vol}(B)$, where $B \subset \langle C \rangle$ is the unit ball centered at the origin. Let F be a k -dimensional face of some solid cone C . By C/F we denote the orthogonal projection of C to the subspace $\langle F \rangle^\perp$. We note that $(C/F)^*$ is an $(n - k)$ -dimensional face of C^* that is orthogonal to F .

Let now $W \subset O(V)$ be a finite reflection group. Set $W^k = \{w \in W : \dim \ker(1 - w) = k\}$ and denote $W^{\text{reg}} := W^0$. For a subspace $U \subset V$ we denote by W_U the subgroup of W that fixes U pointwise.

The aim of this paper is to prove the following theorem conjectured by the first author in [1].

Theorem 1. *For a fundamental cone C of a finite reflection group W and for each $k = 0, \dots, n$ we have*

$$\sum_{F \subset C, \dim F = k} \sigma(F) \cdot \sigma((C/F)^*) = |W^k| / |W|,$$

where F runs over the k -dimensional faces of C .

The following two results are crucial for the proof of Theorem 1. The first one is the fundamental result of Waldspurger [7], for the simplest proof of which we refer the reader to [2, 3]. The second one is the so-called ‘‘Curious Identity’’ of De Concini and Procesi [5] (see also [1, 6]).

Theorem 2 (Waldspurger). $C^* = \bigsqcup_{w \in W} (1 - w)C^\circ$.

Theorem 3 (Curious Identity). $\sigma(C^*) = |W^{\text{reg}}| / |W|$.

Proof of Theorem 3. Theorem 2 implies that $wC^* = \bigcup_{w' \in W^{\text{reg}}} (1 - w')wC$. Thus we obtain

$$|W|\sigma(C^*) = \sum_{w \in W} \sigma(wC^*) = \sum_{w' \in W^{\text{reg}}} \sum_{w \in W} \sigma((1 - w')wC) = \sum_{w' \in W^{\text{reg}}} \sigma((1 - w')V) = |W^{\text{reg}}|.$$

□

Remark. It is easy to see from the previous proof that the number of cones wC^* covering a generic point of V is equal to $|W^{\text{reg}}|$.

Proof of Theorem 1. Consider the sum $S := \sum_{(C, F)} \sigma(F) \cdot \sigma((C/F)^*)$ over all pairs (C, F) , where C is a fundamental cone of W and F is a k -dimensional face of C . This sum is equal to the left hand side of the required formula multiplied by $|W|$. We shall calculate S in a different way.

Let us recall that any two fundamental cones of W with a common face F are conjugate by a unique element of the reflection subgroup W_F that fixes F pointwise. We also note that C/F is a fundamental cone for the action of W_F on $\langle F \rangle^\perp$. By Theorem 3 we have $\sigma((C/F)^*) = |W_F^{\text{reg}}| / |W_F|$. If we take the sum of $\sigma((C/F)^*)$ over all C that contain a fixed face F , we get $|W_F^{\text{reg}}|$. The group W_F and the measure $\sigma((C/F)^*)$ depend only on the subspace $U := \langle F \rangle$ that is an intersection of reflection hyperplanes. Let us take the sum of $\sigma(F) \cdot \sigma((C/F)^*)$ over all pairs (C, F) such that $F \subset U$ for a fixed k -dimensional space U . We get $|W_U^{\text{reg}}|$ multiplied by the total measure of faces $F \subset U$, which is equal to one, since the faces F decompose U . Taking the sum over all subspaces U we obtain $S = |W^k|$. □

¹The first author was partially supported by the Moebius Contest Foundation for Young Scientists

²The second author was partially supported by the following grants RFBR 09-01-00287, 09-01-12169.

Remarks. 1. Given a cone C and its face F , we define the cone $F \oplus (C/F)^*$. It follows from the previous proof that for any k -dimensional subspace U , which is an intersection of reflection hyperplanes, a generic point of V is covered by $|W_U^{\text{reg}}|$ cones $F \oplus (C/F)^*$ with $F \subset U$.

2. The sums from Theorem 1 can be expressed in terms of the exponents $m_1, \dots, m_n$ of W with a help of the Solomon formula [4]: $\sum_{k=0}^n |W^{n-k}| t^k = \prod_{k=0}^n (1 + m_k t)$.

Definition. Let C be a fundamental cone of W . We say that two k -dimensional faces F and F' of C are equivalent if there exists an element $w \in W$ such that $w\langle F \rangle = \langle F' \rangle$.

Denote by N_F the subgroup of W normalizing $\langle F \rangle$. Consider the sum $(|W|/|W_F|) \cdot \sum_{F' \sim F} \sigma(F')$ of relative angle measures for the W -translates of the k -dimensional faces $F' \subset C$ that are equivalent to F . It is the same as the total measure of the W -translates of $F' \subset \langle F \rangle$. The latter is equal to $|W|/|N_F|$ multiplied by the total measure of the faces $F' \subset \langle F \rangle$ which is equal to 1. Thus we get $\sum_{F' \sim F} \sigma(F') = |W_F|/|N_F|$. Using Theorems 1, 3 and applying the previous equality we get:

$$\sum_F |W_F^{\text{reg}}|/|N_F| = \sum_F \left(\sum_{F' \sim F} \sigma(F') \right) \cdot \sigma((C/F)^*) = |W^k|/|W|,$$

where the first sum is taken over representatives of all equivalence classes of k -dimensional faces $F \subset C$.

The authors are grateful to E. B. Vinberg for valuable discussions.

References

- [1] Bibikov P. V. *Relative Angle Measure of the Dual Cone of the Fundamental Cone of a Finite Reflection Group*, Uspehi Mat. Nauk, (2009).
- [2] Bibikov P. V., Zhgoon V. S. *On the Waldspurger theorem*, Uspehi Mat. Nauk, 64:5 (389) (2009), 177–178.
- [3] Bibikov P. V., Zhgoon V. S. *On Tilings Defined by Discrete Reflection groups*, Izvestiya RAN. Ser. Mat., (2009).
- [4] Solomon L. *Invariants of finite reflection groups*, Nagoya Math. J. 22 (1963) 57–64.
- [5] De Concini C., Procesi C. *A curious identity and the volume of the root spherical simplex*, with appendix by J. Stembridge, Atti Accad. Naz. Lincei Cl. Sci. Fis. Mat. Natur. Rend. Lincei (9) Mat. Appl. 17 (2006), no. 2, 155–165.
- [6] Denham G. *A note of De Concini and Procesi's curious identity*, Mat. Appl. v.19 (2008), n.1, 55–63.
- [7] Waldspurger J.-L. *Une remarque sur les systèmes de racines*, Journal of Lie Theory, Volume 17 (2007), Number 3. P. 597–603.